

**Selected Articles on the Analysis of Drugs of Abuse in Seized Materials
(January - June 2020)**

Prodrugs of new psychoactive substances (NPS): A new challenge, S.P. Elliot; T. Holdbrook; S.D.

Brandt, *Journal of Forensic Sciences*, **2020**, 65, 913-920. <https://doi.org/10.1111/1556-4029.14268>

Report on a novel emerging class of highly potent benzimidazole NPS opioids: Chemical and in vitro functional characterization of isotonitazene, P. Blanckaert; A. Cannaert; K. Van Uytvanghe; F. Hulpia; E. Deconinck; S. Van Calenbergh; C. Stove, *Drug Testing and Analysis*, **2020**, 12, 422– 430.

<https://doi.org/10.1002/dta.2738>

Hitting the jackpot – development of gas chromatography–mass spectrometry (GC–MS) and other rapid screening methods for the analysis of 18 fentanyl-derived synthetic opioids, N. Gilbert; L. H. Antonides; C.J. Schofield; A. Costello; B. Kilkelly; A.R. Cain; P.R.V. Dalziel; K. Horner; R.E. Mewis; O.B. Sutcliffe, *Drug Testing and Analysis*, **2020**, 12, 798– 811. <https://doi.org/10.1002/dta.2771>

New psychoactive substances (NPS) prevalence over LSD in blotter seized in State of Santa Catarina, Brazil: A six-year retrospective study, B. de Souza Boff; J.S. Filho; K. Nonemacher; S. Driessen Schroeder M. Dutra Arbo; K. Zomer Rezin, *Forensic Science International*, **2020**, 306, 110002.

<https://doi.org/10.1016/j.forsciint.2019.110002>

Triple quadrupole–mass spectrometry protocols for the analysis of NBOMes and NBOHs in blotter papers, D.R. de Moraes; K.F. da Cunha; T. Betoni Rodrigues; R. Lanaro; L. de Melo Barbosa; J.J. Zacca; M. Nogueira Eberlin; J.L. Costa, *Forensic Science International*, **2020**, 309, 110184.

<https://doi.org/10.1016/j.forsciint.2020.110184>

Identification of new NBOH drugs in seized blotter papers: 25B-NBOH, 25C-NBOH, and 25E-NBOH, Y. Machado; J. Coelho Neto; R.A. Lordeiro; R.B. Alves; E. Piccin, *Forensic Toxicology*, **2020**, 38, 203–215.

<https://doi.org/10.1007/s11419-019-00509-7>

Return of the lysergamides. Part VI: Analytical and behavioural characterization of 1-cyclopropanoyl-*d*-lysergic acid diethylamide (1CP-LSD), S.D. Brandt; P.V. Kavanagh; F. Westphal; A. Stratford; A.U.

Odland; A.K. Klein; G. Dowling; N.M. Dempster; J. Wallach; T. Passie; A.L. Halberstadt, *Drug Testing and Analysis*, **2020**, 12, 812– 826. <https://doi.org/10.1002/dta.2789>

Detection and quantitation of synthetic cannabinoid receptor agonists in infused papers from prisons in a constantly evolving illicit market, C. Norman; G. Walker; B. McKirdy; C. McDonald; D. Fletcher; L.H. Antonides; O.B. Sutcliffe; N. Nic Daéid; C. McKenzie, *Drug Testing and Analysis*, **2020**, 12, 538– 554. <https://doi.org/10.1002/dta.2767>

The phenomenon of *para*-fluorophenylpiperazine (pFPP) in combination with the synthetic cannabinoid AMB-FUBINACA in seized plant material in New Zealand, C.S. Johnson; C.R. Stansfield; V.R. Hassan; E. Kolbe; H.K. Partington; D.C. Kappatos; R.F. Somerville, *Forensic Science International*, **2020**, 307, 110107. <https://doi.org/10.1016/j.forsciint.2019.110107>

Determination of chiral cathinone in fresh samples of *Catha edulis*, A.M. Dhabbah, *Forensic Science International*, **2020**, 307, 110105. <https://doi.org/10.1016/j.forsciint.2019.110105>

An efficient ambient ionization mass spectrometric approach to detection and quantification of the mescaline content of commonly abused cacti from the *Echinopsis Genus*, C.M. Longo; R.A. Musah, *Journal of Forensic Sciences*, **2020**, 65, 61-66. <https://doi.org/10.1111/1556-4029.14134>

The Italian panorama of cannabis light preparation: Determination of cannabinoids by LC-UV, M. Dei Cas; E. Casagni; A. Saccardo; S. Arnoldi; C. Young; S. Scotti; E. Vieira de Manicor; V. Gambaro; G. Roda, *Forensic Science International*, **2020**, 307, 110113. <https://doi.org/10.1016/j.forsciint.2019.110113>

Synthetic route sourcing of illicit at home cannabidiol (CBD) isomerization to psychoactive cannabinoids using ion mobility-coupled-LC-MS/MS, T.D. Kiselak; R. Koerber; G.F. Verbeck, *Forensic Science International*, **2020**, 308, 110173. <https://doi.org/10.1016/j.forsciint.2020.110173>.

Cannabidiol and tetrahydrocannabinol concentrations in commercially available CBD E-liquids in Switzerland, K.E. Grafinger; S. Krönert; A. Broillet; W. Weinmann, *Forensic Science International*, **2020**, 310, 110261. <https://doi.org/10.1016/j.forsciint.2020.110261>

Validation of an UHPLC/DAD method for the determination of cannabinoids in seized materials: Analysis of 213 samples sold in Belgian CBD shops, M. Deville; N. Dubois; R. Denooz; C. Charlier, *Forensic Science International*, **2020**, 310, 110234. <https://doi.org/10.1016/j.forsciint.2020.110234>

Concentrations of THC, CBD, and CBN in commercial hemp seeds and hempseed oil sold in Korea, E. Jang; H. Kim; S. Jang; J. Lee; S. Baeck; S. In; E. Kim; Y. Kim; E. Han, *Forensic Science International*, **2020**, 306, 110064. <https://doi.org/10.1016/j.forsciint.2019.110064>

Identification and characterization of chemically masked derivatives of pseudoephedrine, ephedrine, methamphetamine, and MDMA, A. Mayer; B. Copp; B. Bogun; G. Miskelly, *Drug Testing and Analysis*, **2020**, 12, 524–537. <https://doi.org/10.1002/dta.2764>

Unexpected identification and characterization of a cathinone precursor in the new psychoactive substance market: 3',4'-methylenedioxy-2,2-dibromobutyrophenone, S. Armenta; C. Gil; M. Ventura; F.A. Esteve-Turrillas, *Forensic Science International*, **2020**, 306, 110043. <https://doi.org/10.1016/j.forsciint.2019.110043>

Chemometrics in forensic chemistry — Part II: Standardized applications – Three examples involving illicit drugs, T. Salonen; B. Ahrens; M. Bovens; J. Eliaerts; S. Huhtala; A. Nordgaard; I. Aberink, *Forensic Science International*, **2020**, 307, 110138. <https://doi.org/10.1016/j.forsciint.2019.110138>

Profiling

Organic impurity profiling of 3,4-methylenedioxymethamphetamine (MDMA) synthesised from catechol and eugenol via 4-allylcatechol, E. Heather; R. Shimmon; A. McDonagh, *Forensic Science International*, **2020**, 309, 110176. <https://doi.org/10.1016/j.forsciint.2020.110176>

Identification of specific markers for amphetamines synthesized from glycidic acid pre-precursors and retrospective search in German profiling database, F.M. Hauser; M. Pütz; T. Rößler; J.W. Hulshof, *Drug Testing and Analysis*, **2020**, 12, 41-52. <https://doi.org/10.1002/dta.2686>

Evaluation of data preprocessings for the comparison of GC–MS chemical profiles of seized cannabis samples , A. Slosse; F. Van Durme; N. Samyn; D. Mangelings; Y. Vander Heyden, *Forensic Science International*, **2020**, 310, 110228. <https://doi.org/10.1016/j.forsciint.2020.110228>

Isotopomics by isotope ratio monitoring by ¹³C nuclear magnetic resonance spectrometry on cutting agents in heroin: A new approach for illicit drugs trafficking route elucidation, V. Joubert; M. Trébuchet; M. Mikic; V. Silvestre; A.M. Schiphorst; D. Loquet; A. Stemmelen; V. Ladroue; F. Besacier; S. Akoka; G.S. Remaud, *Drug Testing and Analysis*, **2020**, 12, 449– 457. <https://doi.org/10.1002/dta.2745>

**Selected Articles on the Analysis of Drugs of Abuse in Biological Specimens
(January - June 2020)**

Determination of morphine, fentanyl and their metabolites in small sample volumes using liquid chromatography tandem mass spectrometry, J. Gleba; J. Kim, *Journal of Analytical Toxicology*, **2020**, 44, 325–330. <https://doi.org/10.1093/jat/bkz104>

Toxicokinetic studies and analytical toxicology of the new synthetic opioids cyclopentanoyl-fentanyl and tetrahydrofuranoyl-fentanyl, T.M. Gampfer; L. Wagmann; M.J. Richter; S. Fischmann; F. Westphal; M.R. Meyer, *Journal of Analytical Toxicology*, **2020**, 44, 449–460. <https://doi.org/10.1093/jat/bkaa010>

Development of an UPLC–MS/MS method for the analysis of 16 synthetic opioids in segmented hair, and evaluation of the polydrug history in fentanyl analogue users, M. del Mar Ramirez Fernandez; S.M.R. Wille; D. Jankowski; V. Hill; N. Samyn, *Forensic Science International*, **2020**, 307, 110137. <https://doi.org/10.1016/j.forsciint.2019.110137>

Effects of lipophilicity and functional groups of synthetic cannabinoids on their blood concentrations and urinary excretion, H. Kakeashi; N. Shima; A. Ishikawa; A. Nitta; R. Asai; M. Wada; S. Nakano; S. Matsuta; K. Sasaki; H. Kamata; T. Kamata; H. Nishioka; A. Miki; M. Katagi, *Forensic Science International*, **2020**, 307, 110106. <https://doi.org/10.1016/j.forsciint.2019.110106>

Emerging synthetic cannabinoids: development and validation of a novel liquid chromatography quadrupole time-of-flight mass spectrometry assay for real-time detection, A.J. Krotulski; A.L.A. Mohr; B.K. Logan, *Journal of Analytical Toxicology*, **2020**, 44, 207–217. <https://doi.org/10.1093/jat/bkz084>

Solid-phase extraction–liquid chromatography–tandem mass spectrometry method for the qualitative analysis of 61 synthetic cannabinoid metabolites in urine, F. Gaunitz; T. Kieliba; M. Thevis; K. Mercer-Chalmers-Bender, *Drug Testing and Analysis*, **2020**, 12, 27-40. <https://doi.org/10.1002/dta.2680>

Simultaneous analysis of 29 synthetic cannabinoids and metabolites, amphetamines, and cannabinoids in human whole blood by liquid chromatography–tandem mass spectrometry – A New Zealand perspective of use in 2018, R.S. Ong; D.C. Kappatos; S.G.G. Russell; H.A. Poulsen; S.D. Banister; R.R. Gerona; M. Glass; C.S. Johnson; M.J. McCarthy, *Drug testing and Analysis*, **2020**, 12, 195-214. <https://doi.org/10.1002/dta.2697>

5F-MDMB-PICA metabolite identification and cannabinoid receptor activity, M.T. Truver; S. Watanabe; A. Åstrand; S. Vikingsson; H. Green; M.J. Swortwood; R. Kronstrand, *Drug Testing and Analysis*, **2020**, 12, 127-135. <https://doi.org/10.1002/dta.2688>

Detection and characterization of the new synthetic cannabinoid APP-BINACA in forensic casework, A.J. Krotulski; A.L.A. Mohr; F.X. Diamond; B.K. Logan, *Drug Testing and Analysis*, **2020**, 12, 136-144. <https://doi.org/10.1002/dta.2698>

Forty-three fatalities involving the synthetic cannabinoid, 5-fluoro-ADB: Forensic Pathology and Toxicology Implications, D.M. Boland; L.J. Reidy; J.M. Seither; J.M. Radtke; E.O. Lew, *Journal of Forensic Sciences*, **2020**, 65, 170-182. <https://doi.org/10.1111/1556-4029.14098>

Simultaneous determination of synthetic cannabinoids and their metabolites in human hair using LC-MS/MS and application to human hair, B. Cho; H.S. Cho; J. Kim; J. Sim; I. Seol; S.K. Baeck; S. In; D.H. Shin; E. Kim, *Forensic Science International*, **2020**, 306, 110058. <https://doi.org/10.1016/j.forsciint.2019.110058>

Solid-phase extraction based on cation-exchange sorbents followed by liquid chromatography high-resolution mass spectrometry to determine synthetic cathinones in urine, S. Pascual-Caro; N. Fontanals; F. Borrull; C. Aguilar; M. Callul, *Forensic Toxicology*, **2020**, 38, 185-194. <https://doi.org/10.1007/s11419-019-00508-8>

Development and validation of a GC-MS/MS method for the determination of 11 amphetamines and 34 synthetic cathinones in whole blood, M.K. Woźniak; L. Banaszkiwicz; M. Wiergowski; E. Tomczak; M. Kata; B. Szpiech; J. Namiesnik; M. Biziuk, *Forensic Toxicology*, **2020**, 38, 42-58. <https://doi.org/10.1007/s11419-019-00485-y>

Case report on two-cathinones abuse: MPHP and *N*-ethyl-4'-methylnorpentadone, with a fatal outcome, B. Lelievre; C. Richeval; A. Coulon; D. Iwanikow; M. Brofferio; M. Deguigne; D. Boels; D. Allorge; S. Ferec; G. Drevin; N. Jousset; J.M. Gaulier, *Forensic Toxicology*, **2020**, 38, 243-254. <https://doi.org/10.1007/s11419-019-00486-x>

Studies on the phase I metabolites of the new designer drug 1-(2,3-dihydro-1*H*-inden-5-yl)-2-(pyrrolidine-1-yl)butan-1-one (5-PPDI) in human urine, A. Ishii; K. Kusakabe; N. Kato; S. Sasaki; K. Tsujikawa; T. Wada, *Forensic Science International*, **2020**, 310, 110214. <https://doi.org/10.1016/j.forsciint.2020.110214>

Fatal intoxication with *N*-ethylpentylone: a case report and method for determining *N*-ethylpentylone in biological material, M. Zawadzki; K. Nowak; P. Szpot, *Forensic Toxicology*, **2020**, 38, 255-263. <https://doi.org/10.1007/s11419-019-00483-0>

Structural elucidation of phenidate analogues via the ESI-MS/MS spectra of their sodium adduct ions, T.S. Yamin; H. Prihed; M. Madmon; A. Shifrovitch; A. Bratz; A. Weissberg, *Forensic Science International*, **2020**, 306, 110044. <https://doi.org/10.1016/j.forsciint.2019.110044>

Simultaneous quantitative determination of amphetamines, opiates, ketamine, cocaine and metabolites in human hair: Application to forensic cases of drug abuse, Y. Zhuo; X. Wang; J. Wu; S. Zhang; H. Deng; W. Liu; H. Wu; P. Xiang; M. Shen, *Journal of Forensic Sciences*, **2020**, 65, 563-569. <https://doi.org/10.1111/1556-4029.14179>

An evaluation of the cut-off value of methamphetamine in hair samples via HPLC-MS/MS, J. Ou; Y. Zhang; S. Chen; C. Zhong; Q. Xu; D. Xu; X. Yan, *Forensic Science International*, **2020**, 306, 110094. <https://doi.org/10.1016/j.forsciint.2019.110094>

Comparison of the detection windows of heroin metabolites in human urine using online SPE and LC-MS/MS: Importance of morphine-3-glucuronide, L. Wang; C. Ni; H. Shen; Z. Sheng; C. Liang; R. Wang; Y. Zhang, *Journal of Analytical Toxicology*, **2020**, 44, 22-28. <https://doi.org/10.1093/jat/bkz040>

Comparative analysis of ELISA immunoassay and LC-QTOF for opiate screening, D. Kennedy; M. Dani, *Journal of Analytical Toxicology*, **2020**, 44, 410-413. <https://doi.org/10.1093/jat/bkz109>

Pitfalls in drug testing by hyphenated low- and high-resolution mass spectrometry, H.H. Maurer, *Drug Testing and Analysis*, **2020**, 12, 172-179. <https://doi.org/10.1002/dta.2744>