

8. REFERENCES

Panda S, Mallick PS, Karim MA, Sharifuzzaman M, Ahmed AHT, Baatsen P, 2002, National assessment of situation and responses to opiod/opiate use in Bangladesh (NASROB). Dhaka, FHI/CARE/HASAB.

Panda S, Bijaya L, Sadhana Devi N, Foley E, Chatterjee A, Banerjee D, Naik TN, Saha MK and Bhattacharya S, 2001, 'Interface between drug use and sex work in Manipur'. *National Medical Journal of India*, 14: 209 – 211.

Panda S, Chatterjee A, Bhattacharya SK, Manna B, Singh PN, Sarkar S, Naik TN, Chakrabarti S, Detels R, 2000, 'Transmission of HIV from injecting drug users to their wives in India'. *International Journal of STD & AIDS*, 11: 468 – 473.

UNODC and UNAIDS, 1999, 'Baseline study of the relationship between injecting drug use, HIV and Hepatitis C among male injecting drug users in Lahore'. UNODCCP/UNAIDS Pakistan, December.

Further reading on assessment guides and estimation techniques

FHI-Impact, USAID, UNAIDS, WHO and UNDCP, 2003, 'Estimating the size of populations at risk for HIV: issues and methods'.

International HIV/AIDS Alliance, 2003, 'Developing HIV/AIDS work with drug users: a guide to participatory assessment and response'.

Mental health: research and evidence, Department of mental health and substance dependence. Non-communicable Disease and Mental Health Cluster, World Health Organization, 2002 'Sex-RAR guide: the rapid assessment and response guide on psychoactive substance use and sexual risk behavior'.

UNODCCP, 1999, 'Drug abuse rapid situation assessments and responses'.

World Health Organization and United Nations International Children's Education Fund, 1998, The rapid assessment and response guide on psychoactive substance abuse and especially vulnerable young people. Geneva: WHO and UNICEF.

World Health Organization and United Nations Joint Programme on HIV/AIDS, 1998, 'The rapid assessment and response guide on substance use and sexual risk behavior'. Geneva: WHO and UNAIDS.

World Health Organization, 1998, The rapid assessment and response guide on injecting drug use (draft for field testing). Geneva: WHO, 1998.