



Global seizures of ecstasy-type substances^(a): 2006-2016




Year	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
Kg equivalents ^(b)	9,782	18,839	7,510	6,558	6,074	4,424	5,296	4,917	9,601	10,191	13,963

SEIZURES OF ECSTASY-TYPE SUBSTANCES^(a) - as percentage of world total and in kg equivalents^(b) - HIGHEST RANKING COUNTRIES - 2016



SEIZURES OF ECSTASY-TYPE SUBSTANCES BY REGION - as percentage of world total and in kg equivalents^(a) - 2016




^(a) Aside from seizures of ecstasy-type substances (e.g. MDMA, MDA, MDE), includes substances reported as "ecstasy" which may not have been confirmed by forensic testing.


^(b) This quantity reflects the bulk weight of seizures, with no adjustment for purity. Seizures reported by volume (litres, millilitres, etc.) are converted to kg equivalents by assuming a weight of 1 kg per litre. Seizures of ecstasy-type substances in tablets or similar units are converted using assumed bulk tablet weights between 200mg and 300 mg, depending on the region and specific drug type, and based on information currently available to UNODC. Further details on conversion factors are given in the methodology section accompanying the World Drug Report.

^(c) Data from England & Wales only. Financial year 2015/16


**ECSTASY-TYPE SUBSTANCES*
INTERCEPTED - WORLD: 2006 - 2016**




**ECSTASY-TYPE SUBSTANCES*
INTERCEPTED - ASIA: 2006 - 2016**




ECSTASY-TYPE SUBSTANCES* INTERCEPTED - AMERICAS: 2006 - 2016




**ECSTASY-TYPE SUBSTANCES*
INTERCEPTED - EUROPE: 2006 - 2016**



**ECSTASY-TYPE SUBSTANCES*
INTERCEPTED - AFRICA: 2006 - 2016**



**ECSTASY-TYPE SUBSTANCES*
INTERCEPTED - OCEANIA: 2006 - 2016**



*Aside from seizures of ecstasy-type substances (e.g. MDMA, MDA, MDE), includes substances reported as "ecstasy" which may not have been confirmed by forensic testing.

** This quantity reflects the bulk weight of seizures, with no adjustment for purity. Seizures reported by volume (litres, millilitres, etc.) are converted to kg equivalents by assuming a weight of 1 kg per litre. Seizures of ecstasy-type substances reported in tablets or similar units are converted using assumed bulk tablet weights between 200mg and 300 mg, depending on the region and specific drug type, and based on information currently available to UNODC. Further details on conversion factors are given in the methodology section accompanying the World Drug Report.